#### IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

PATENT

### DIVISIONAL APPLICATION REQUEST TRANSMITTAL

SISTANT COMMISSIONER OF PATENTS

ashington, D.C. 20231

TIP.

Attorney Docket No. XI/P03139US2
Prior Application Information:
Art Unit: 1615
Examiner: N. Levy
Serial No.: 08/863,182

Transmitted herewith for filing is a Rule 53(b) Divisional patent application

of: JEANNIN, Philippe

for: N-PHENYLPYRAZOLE-BASED ANTI-FLEA AND ANTI-TICK COLLAR FOR CATS AND DOGS.

#### ENCLOSED are the following:

- 1. [XX] Specification, abstract and claims of 32 pages.
- 2. [XX] sheets of [ ] formal [ ] informal drawings [ XX ] No drawings.
- 3. [XX] Declaration (original or copy) signed by the named inventor.
- 4. [XX] Preliminary Amendment
- I Information Disclosure Statement
- 6. [ ] Other:

#### NOTE the following:

- [ ] Applicant is a small entity, which was established in parent application Serial No. \*, and such status is still proper and desired.
- 8. [XX] The prior application is assigned to RHONE MERIEUX.
- 9. [XX] Priority of the following applications are claimed under 35 U.S.C. 119:

		*	Certified Copy	of Priority Doc. Filed
Serial No.	Date Filed	Country	in USSN	Date
96 04 206	29 March 1996	France	08/863,182	May 27, 1997
96 03 707	26 March 1997	France	08/863,182	December 17, 1997

- 10. [XX] Amend the specification by inserting before the first line the sentence:
  - a. [ ] This is a continuation of application Serial No. \* filed \*.
  - b. [XX] This is a divisional of application Serial No. 08/863,182 filed May 27, 1997; which is a CIP of 08/692,430 filed 08/05/96 now abandoned.



11. [ ] Cancel claims prior to calculation of the filing fee.

FILING FEE: calculated below (after reduction for amendment or claims cancellations if noted above):

[X] Total claims:	50	_	20	===	30	х	\$ 22.00	=	\$ 660.00
[X] Independent claims:	3	-	3	=	0	x	\$ 82.00	=	\$
[ ] Multiple dependent claim(s	(					+	\$270.00	==	\$
						BA	SIC FEE	=	\$ 790.00
TOTAL OF ABOVE CALCULATIONS									\$
[] Reduction by 1/2 for filing	by s	mal	l entit	ty				_	<u>\$[1</u>
						SU	BTOTAL	=	\$1,450.00
[ ] Fee for extension of time in parent application									\$
TOTAL OF FEES								=	\$1,450,00

- [XX] A check in the amount of \$1,450.00 is enclosed (must at least cover the basic fee). If no check or an insufficient check is enclosed and a fee is due herewith, the Commissioner is authorized to charge any fee or additional fee due in connection herewith to Deposit Account No. 12-0555. A duplicate of this sheet is enclosed.
- [XX] In the event that a petition for extension of time is required to be submitted in the pending prior application, and in the event that a separate petition for the pending prior application is not filed on even date, applicant hereby petitions under 37 C.F.R. §1.136(a) for an extension of time in the pending prior application of as many months as are required to render this submission timely. The Commissioner is authorized to charge any fee or additional fee due in connection with this communication to Deposit Account No. 12-0555. A duplicate copy of this sheet is enclosed.

Respectfully submitted

Date: October 19, 1998

Thomas P. Sarro
Registration No. 19,396

LARSON & TAYLOR • Suite 900 • 1199 North Fairfax St. • Alexandria, VA 22314

# Rule 53(b) Divisional Application Filed Concurrently Herewith

# IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of:

Attorney's Docket: XI/P03139US2

JEANNIN, Philippe

Divisional Application of

Serial No. 08/863,182

: Art Unit: 1615

Filed: Concurrently herewith

: Examiner: N. Levy

For: N-PHENYLPYRAZOLE-BASED ANTI-FLEA AND ANTI-TICK COLLAR FOR CATS

AND DOGS

#### PRELIMINARY AMENDMENT

The Assistant Commissioner of Patents Washington, D.C.

SIR:

Preliminary to examination of the above-identified application, please amend the application as follows:

## IN THE CLAIMS

Please amend claim 1 as follows:

1. (Amended) Anti-flea and anti-tick [collar or other] external device for a pet, [in particular a cat or dog, made of] to ensure more than six months of efficacy of greater than 95% against fleas, as determined in a test providing the reinfestation of the pet with

100 \* 10 fleas, and more than three months of efficacy of greater than 90% against ticks, as determined in a test providing the reinfestation of the pet with 50 \* 3 three ticks, comprising a matrix in which is incorporated from 0.1 to 40% by weight, relative to the [collar] external device, of a substance which is active against fleas and ticks, this active substance being formed of at least one compound corresponding to formula (I) below

in which:

R<sub>1</sub> is CN or methyl or a halogen atom;

R<sub>2</sub> is S(O)<sub>n</sub>R<sub>3</sub> or 4,5-dicyanoimidazol-2-yl or haloalkyl;

R<sub>3</sub> is alkyl or haloalkyl;

 $R_4$  represents a hydrogen or halogen atom; or a radical  $NR_5R_5$ ,  $S(O)_mR_7$ ,  $C(O)R_7$ ,  $C(O)O-R_7$ , alkyl, haloalkyl or  $OR_8$  or a radical -N=C( $R_9$ ) ( $R_{10}$ );

 $R_s$  and  $R_s$  independently represent a hydrogen atom or an alkyl, haloalkyl, C(O)alkyl, alkoxycarbonyl or S(O),CF $_3$  radical; or  $R_s$  and  $R_s$  may together form a divalent alkylene radical which may be interrupted by one or two divalent hetero atoms, such as oxygen or sulphur;

R, represents an alkyl or haloalkyl radical;

R<sub>s</sub> represents an alkyl or haloalkyl radical or a hydrogen atom;

Ro represents an alkyl radical or a hydrogen atom;

R<sub>10</sub> represents a phenyl or heteroaryl group optionally substituted with one or more halogen atoms or groups such an OH, -O-alkyl, -S-alkyl, cyano or alkyl;

 $\rm R_{11}$  and  $\rm R_{12}$  represent, independently of each other, a hydrogen or halogen atom, or CN or NO $_{\circ}$ ;

 $\rm R_{13}$  represents a halogen atom or a haloalkyl, haloalkoxy,  $\rm S(O)_q \rm CF_3$  or  $\rm SF_5$  group;

m, n, q and r represent, independently of each other, an integer equal to 0, 1 or 2;

X represents a trivalent nitrogen atom or a radical C-R<sub>12</sub>, the other three valency positions of the carbon atom forming part of the aromatic ring;

with the proviso that when  $R_1$  is methyl, either  $R_3$  is haloalkyl,  $R_4$  is  $NH_2$ ,  $R_{11}$  is Cl,  $R_{13}$  is  $CF_3$  and X is N; or  $R_2$  is 4,5-dicyanoimidazol-2-yl,  $R_4$  is Cl,  $R_{11}$  is CL,  $R_{13}$  is  $CF_3$ , and X is =C-Cl[:]

[this collar or other external device being designed to ensure more than 6 months of efficacy against fleas and more than 3 months of efficacy against ticks, the efficacy preferably being maintained for several weeks even if the collar or other external device is taken off or lost or if there is a variation in the release of the compound (I) by the matrix].

Please add the following new claims 48-52:

--48. Anti-flea and anti-tick device according to claim 1 wherein the device ensures more than six months of efficacy against fleas and more than three months of efficacy against ticks.

49. Method for eliminating fleas and ticks from cats and dogs comprising: attaching to the cats and dogs at least one external device which comprises a matrix incorporating from 0.1 to 40% by weight, relative to the [collar] external device, of a substance which is active against fleas and ticks, this active substance being formed of at least one compound corresponding to formula (I) below

in which:

R<sub>1</sub> is CN or methyl or a halogen atom;

 $R_2$  is  $S(O)_nR_3$  or 4,5-dicyanoimidazol-2-yl or haloalkyl;

R<sub>3</sub> is alkyl or haloalkyl;

 $R_4 \ \text{represents a hydrogen or halogen atom; or a radical } NR_5R_6, \ S(O)_mR_7, \\ C(O)R_7, \ C(O)O-R_7, \ \text{alkyl}, \ \text{haloalkyl or } OR_6 \ \text{or a radical -N=C}(R_9) \ (R_{10});$ 

 $R_{\rm 5}$  and  $R_{\rm 6}$  independently represent a hydrogen atom or an alkyl, haloalkyl, C(O)alkyl, alkoxycarbonyl or S(O),CF $_{\rm 3}$  radical; or  $R_{\rm 6}$  and  $R_{\rm 6}$  may together form a divalent alkylene radical which may be interrupted by one or two divalent hetero atoms, such as oxygen or sulphur;

R<sub>7</sub> represents an alkyl or haloalkyl radical;

R<sub>8</sub> represents an alkyl or haloalkyl radical or a hydrogen atom;

R<sub>9</sub> represents an alkyl radical or a hydrogen atom;

 $R_{10}$  represents a phenyl or heteroaryl group optionally substituted with one or more halogen atoms or groups such an OH, -O-alkyl, -S-alkyl, cyano or alkyl:

 $R_{\rm 11}$  and  $R_{\rm 12}$  represent, independently of each other, a hydrogen or halogen atom, or CN or  $NO_2;$ 

 $\rm R_{13}$  represents a halogen atom or a haloalkyl, haloalkoxy,  $\rm S(O)_q \rm CF_3$  or  $\rm SF_5$  group;

m, n, q and r represent, independently of each other, an integer equal to 0, 1 or 2;

X represents a trivalent nitrogen atom or a radical C-R<sub>12</sub>, the other three valency positions of the carbon atom forming part of the aromatic ring;

with the proviso that when  $R_1$  is methyl, either  $R_3$  is haloalkyl,  $R_4$  is  $NH_2$ ,  $R_{11}$  is Cl,  $R_{13}$  is CF $_3$  and X is N; or  $R_2$  is 4,5-dicyanoimidazol-2-yl,  $R_4$  is Cl,  $R_{11}$  is CL,  $R_{13}$  is CF $_3$ , and X is =C-Cl;

allowing external device to remain in place in order for compound (I) to be taken up by the animal's sebum and distributed over the animal's entire body, and to be concentrated in the sebaceous glands which become a reservoir for formula (I),

in such a manner that it is ensures more than six months of efficacy of more than 95% against fleas, as determined in a test providing reinfestation with 100  $^{\pm}$  10 fleas, and more than three months of efficacy of greater than 90% against ticks, as determined in a test providing reinfestation with 50  $^{\pm}$  3 ticks, and that one obtains efficacy during several weeks against ticks and fleas, if the collar or other external is taken off or lost and if there is a variation in the rate of release of the formula (I) by the matrix.

- 50. Method according to claim 49, wherein the external device is left in place for at least 24 hours.
- 51. Method according to claim 49, wherein one obtains an efficacy of two to three months against fleas if the external device is taken off or lost of if there is a variation in the rate of release of the formula (I) by the matrix.
- 52. Method according to claim 49, wherein one obtains an efficacy of one to two months against ticks if the external device is taken off or lost or if there is a variation in the rate of release of the formula (I) by the matrix.—

Please cancel claim 12 on page 26, lines 22 and 23.

Please cancel claims 39 and 41 without prejudice.

Please amend the claims as follows:

Claim 11, line 5, delete "and its derivatives".

21. (Amended) Method for eliminating fleas and ticks from pets to ensure more than six months of efficacy of greater than 95% against fleas, as determined in a test providing the reinfestation of the pet with 100 ± 10 fleas, and more than three months of efficacy of greater than 90% against ticks, as determined in a test providing the reinfestation of the pet with 50 ± 3 three ticks, [in particular cats and dogs, in which one attaches] comprising attaching to the pets at least one [collar or other] external device [which comprises] having a matrix into which is incorporated a compound corresponding to formula (I) below:

$$R_{1}$$
 $R_{11}$ 
 $R_{13}$ 
 $R_{13}$ 

in which:

R, is CN or methyl or a halogen atom;

R<sub>2</sub> is S(O)<sub>n</sub>R<sub>3</sub> or 4,5-dicyanoimidazol-2-yl or haloalkyl;

R<sub>3</sub> is alkyl or haloalkyl;

 $R_4 \text{ represents a hydrogen or halogen atom; or a radical } NR_8R_6, \ S(O)_mR_7, \\ C(O)R_7, \ C(O)O-R_7, \ alkyl, \ haloalkyl \ or \ OR_8 \ or \ a \ radical \ -N=C(R_9) \ (R_{10});$ 

 $R_{\rm s}$  and  $R_{\rm s}$  independently represent a hydrogen atom or an alkyl, haloalkyl, C(O)alkyl, alkoxycarbonyl or S(O),CF $_{\rm 3}$  radical; or  $R_{\rm s}$  and  $R_{\rm s}$  may together form a divalent alkylene radical which may be interrupted by one or two divalent hetero atoms, such as oxygen or sulphur;

R<sub>7</sub> represents an alkyl or haloalkyl radical;

R<sub>s</sub> represents an alkyl or haloalkyl radical or a hydrogen atom;

R<sub>9</sub> represents an alkyl radical or a hydrogen atom;

 $\rm R_{10}$  represents a phenyl or heteroaryl group optionally substituted with one or more halogen atoms or groups such an OH, -O-alkyl, -S-alkyl, cyano or alkyl;

 $\rm R_{11}$  and  $\rm R_{12}$  represent, independently of each other, a hydrogen or halogen atom, or CN or NO $_{\rm si}$ 

 $\rm R_{13} \ represents \ a \ halogen \ atom \ or \ a \ haloalkyl, \ haloalkoxy, \ S(O)_qCF_3 \ or \ SF_5$  group;

m, n, q and r represent, independently of each other, an integer equal to 0, 1 or 2;

X represents a trivalent nitrogen atom or a radical C-R<sub>12</sub>, the other three valency positions of the carbon atom forming part of the aromatic ring;

with the proviso that when  $R_1$  is methyl, either  $R_3$  is haloalkyl,  $R_4$  is  $NH_2$ ,  $R_{11}$  is CI,  $R_{13}$  is  $CF_3$  and X is N; or  $R_2$  is 4,5-dicyanoimidazol-2-yl,  $R_4$  is CI,  $R_{11}$  is CL,  $R_{13}$  is  $CF_3$ , and X is =C-CI[:]

[which method ensuring prevention and treating fleas and ticks to a high degree of efficacy and other a period exceeding 6 months against fleas and 3 months against ticks, the efficacy preferably being maintained over several weeks even if the collar or external device is taken off or if there is a variation in the release of the compound (I) by the collar or the external device].

Claim 31, line 5, delete "and its derivatives.

Claim 32, line 2, delete "proportion" and insert --concentration--.

Claim 33, line 2, delete "proportion" and insert --concentration--.

Claim 34, line 2, delete "proportion" and insert --concentration--.

Claim 35, line 2, delete "proportion" and insert --concentration--.

Claim 36, line 2, delete "proportion" and insert --concentration--.

Claim 37, line 2, delete "proportion" and insert --concentration--.

Claim 38, line 2, delete "proportion" and insert --concentration--.

Claim 44, lines 1-2, delete "long-lasting".

Claim 45, lines 1-2, delete "long-lasting".

Claim 46, lines 1-2, delete "long-lasting".

Claim 47, line 2, delete "collar or".

#### REMARKS

Entry of the above amendment is respectfully requested.

Respectfully submitted,

LARSON AND TAYLOR

Date: October 19, 1998

Thomas P. Sarro / Registration No. 19,396

Suite 900 1199 North Fairfax Street Alexandria, Virginia, 22314 (703) 739-4900

15

20

2.5

30

# N-Phenylpyrazole-based anti-flea and anti-tick collar for cats and dogs

The present invention relates to an external antiparasitic device, in particular a collar, for pets, in particular cats and dogs, this collar being active against the ectoparasites of these animals, in particular fleas and ticks.

The invention also relates to the use of active compounds for the manufacture of such collars or external devices, as well as to a treatment process relating thereto.

The invention is directed mainly towards fleas of the genus Ctenocephalides, in particular C. felis and C. canis, and ticks, in particular of the genus Rhipicephalus, especially sanguineus, as well as harvest ticks (Trombicula automnalis), which are acarids that mainly attack hunting dogs.

Collars intended to eliminate common ectoparasites from cats and dogs have been produced for a long time. These collars consist of a matrix, usually a plastic matrix, which incorporates between 5 and 40% active substance and is capable of releasing it over time. These collars thus theoretically have the aim of ensuring long-lasting protection.

However, despite the activity claims, out in the field the collars do not have the efficacy required to ensure the actual elimination of these parasites. The cause of this may be the low activity of the active substance included in the matrix. Another cause may be

15

25

30

the accelerated degradation of these active substances under the effect of climatic factors, such as light, heat and rain. Lastly, control of the release of the active substance from the matrix is largely overevaluated. Release generally proves to be difficult and variable, and may depend greatly on the manufacturing conditions, which may vary from one batch to another, and on the conditions of use, in particular climatic variations and especially humidity and temperature, etc. In addition, only a relatively small amount of the active substance incorporated is actually released and it proves difficult to be able to control and optimize its release.

Another drawback of the collars encountered in practice arises from the mode of use of this device which may, obviously, be taken off, worn irregularly, or even be pulled off when the animal moves about, for example in undergrowth; the problem is particularly critical for hunting dogs whose collars are removed before a hunting outing even though they will be confronted with a flea- and tick-ridden environment.

Patent applications WO-A-87/03781 and EP-A-0,295,117 have proposed insecticides of the N-phenyl-pyrazole class.

These substances are described as being active against a very large number of parasites encountered in various fields, namely agriculture, public health and human and veterinary medicine. In the latter field, these substances may act in particular against the fleas and ticks of pets, such as cats and dogs. These substances may be applied in various ways, namely via the oral, parenteral, percutaneous or topical route. The latter type of administration itself covers many

10

15

20

25

30

possibilities, namely sprays, powders, baths, showers, jets, greases, shampoos, creams, waxes, preparations of skin solution type (pour-on) and external devices such as earrings and collars to provide local or systemic treatment. EP-A-0,295,117 and EP-A-0,500,209 propose a slow release composition which may be in the form of a collar or earrings for controlling harmful insects. Such a formulation may comprise from -0.5 - to 25% active material, from 75 to 99.5% polyvinyl chloride and a catalytic amount of a plasticizer, dioctyl phthalate.

French patent application FR-A-2,713,889 describes a pesticide composition containing an insect growth regulator and an N-aryldiazole derivative chosen from the derivatives 4-(2-bromo-1,1,2,2-tetrafluoroethyl)-1-(3-chloro-5-trifluoromethylpyridine-2-yl)-2methylimidazole, 5-amino-3-cyano-1-(2,6-dichloro-4trifluoromethylphenyl)-4-trifluoromethylsulphinylpyra-5-amino-3-cyano-1-(2,6-dichloro-4trifluoromethylphenyl)-4-trifluoromethylthiopyrazole. As in the prior documents discussed above, the compositions are directed towards a large number of insects in different fields, and various types of formulations. Dog fleas and cat fleas are targeted. Among the formulations proposed, this document mentions resinous preparations which may be made into pesticide collars for animals.

However, none of these documents describes the use of pesticidal collars comprising a compound of the N-phenylpyrazole family to control fleas and ticks on pets such as cats and dogs, which make it possible to ensure a high level of efficacy for a long period against these parasites.

As for the pesticidal compounds of the prior art, a specialist might have expected to encounter

10

15

20

25

30

conventional problems of release from the collars and thus problems of activity.

After all, French patent application FR-A-2,713,889 proposes, does it not, the combination of such compounds with an insect growth regulator?

The publications C. Genchi et al., Professione Veterinaria No. 1, supplement 1995, pages 19 to 22, J.M. Postal, Professione Veterinaria No. 1, supplement 1995, pages 17 and 18 and A. Searle et al., Australian Veterinary Practitioner, volume 25, No. 3, 1995, pages 157 and 158 propose instead to effectively control fleas and ticks using a sprayable solution (spray) containing this type of active compound.

The Applicant has now found, surprisingly, that despite the fact that these N-phenylpyrazoles encounter the same difficulties of release as the products of the prior art, it is however possible to obtain collars which are entirely effective for the elimination of cat and dog ectoparasites over a very long period, for example from 6 to 18 months, and that, in addition, the efficacy persists long after the collar is taken off, namely over a period which may be equal to or exceed 2 months, such that it is then possible to obtain a collar which is entirely effective irrespective of the conditions of use. On account of the fact that the protection persists after the collar is taken off, it may be understood that irregular or voluntary use or that resulting from loss of the collar does not jeopardize the protection from which the animal should benefit.

In addition, the Applicant has observed that this long-lasting efficacy was obtained with concentrations of active substance in the matrix forming the

15

2.0

25

3.0

collar which were much smaller than those of the standard products. It was also observed that this efficacy was obtained within a very short time of the collars being put on, in particular an efficacy of greater than 95% in 24 h against fleas and greater than 90% in 48 h against ticks.

It was observed, very surprisingly, that the compounds according to the invention, which are very lipophilic and of high vapour pressure (low volatility), had a very high affinity for the sebum which usually covers the animal's coat (skin and hair), such that, when released, this compound is taken up by the sebum, after which a translocation phenomenon occurs ensuring distribution of active substance over the animal's entire body. In addition, and this is a noteworthy point, these active substances become concentrated in the sebaceous glands which become a reservoir for them, ensuring very long-lasting efficacy and making it possible to compensate for the absence of the collar, by releasing the active substance by passive diffusion.

By virtue of this phenomenon, the variations in release of the active substance by the collar on account of, for example, a variation in climatic conditions, are compensated for by the possibilities of release by the sebaceous glands.

It was also observed that after a bath, which could possibly lead to the removal of the sebum distributed over the animal's body, the animal very rapidly became reprotected, in the presence or absence of the collar, owing to the fact that secretion of new sebum is accompanied by a release, by the sebaceous glands, of the active substance that they contain.

The subject of the present invention is thus

a collar or other external device for a pet, in particular a cat or dog, made of a matrix in which is incorporated from 0.1 to 40% by weight, relative to the collar, of a substance which is active against ectoparasites such as fleas and ticks (anti-flea and anti-tick collar or other external device), this active substance being formed of at least one compound corresponding to formula (I) below:

10

5

15

25

in which:

20

 $R_1$  is CN or methyl or a halogen atom;  $R_2 \text{ is } S(O)_n R_3 \text{ or } 4,5\text{-dicyanoimidazol-2-yl or}$ 

haloalkyl;

R, is alkyl or haloalkyl;

 $R_4 \text{ represents a hydrogen or halogen atom; or}$  a radical NR<sub>5</sub>R<sub>6</sub>, S(O) $_{\rm m}$ R<sub>7</sub>, C(O)R<sub>7</sub>, C(O)O-R<sub>7</sub>, alkyl, haloal-kyl or OR<sub>8</sub> or a radical -N=C(R<sub>9</sub>)(R<sub>10</sub>);

 $R_{5}$  and  $R_{6}$  independently represent a hydrogen atom or an alkyl, haloalkyl, C(O)alkyl, alkoxycarbonyl or S(O) $_{2}\mathrm{CF}_{3}$  radical; or  $R_{5}$  and  $R_{6}$  may together form a divalent alkylene radical which may be interrupted by one or two divalent hetero atoms, such as oxygen or sulphur;

R, represents an alkyl or haloalkyl radical;

10

20

25

30

 $$R_{\vartheta}$$  represents an alkyl or haloalkyl radical or a hydrogen atom;

 $$\rm R_{9}$$  represents an alkyl radical or a hydrogen atom;

R<sub>10</sub> represents a phenyl or heteroaryl group optionally substituted with one or more halogen atoms or groups such as OH, -O-alkyl, -S-alkyl, cyano or alkyl;

 $$R_{11}$$  and  $$R_{12}$$  represent, independently of each other, a hydrogen or halogen atom, or optionally CN or NO,;

 $$R_{13}$$  represents a halogen atom or a haloalkyl, haloalkoxy,  $S(0)\,_{q}CF_{3}$  or  $SF_{5}$  group;

15 . X represents a trivalent nitrogen atom or a radical  $C-R_{12}$ , the other three valency positions of the carbon atom forming part of the aromatic ring;

with the proviso that when  $R_1$  is methyl, either  $R_3$  is haloalkyl,  $R_4$  is  $NH_2$ ,  $R_{11}$  is C1,  $R_{13}$  is  $CF_3$  and X is N; or  $R_2$  is 4,5-dicyanoimidazol-2-yl,  $R_4$  is C1,  $R_{11}$  is C1,  $R_{13}$  is  $CF_3$  and X is =C-C1;

this collar or other external device being designed to ensure at least more than 6 months of efficacy against fleas and at least more than 3 months of efficacy against ticks, the efficacy preferably being maintained for several weeks even if the collar or other external device is taken off or lost or if there is a variation in the rate of release of the compound (I) by the matrix.

Preferably, in formula (I),
R<sub>1</sub> is CN or methyl;
R<sub>2</sub> is S(O)<sub>n</sub>R<sub>3</sub>;
R<sub>3</sub> is alkyl or haloalkyl;

10

15

20

25

30

 $$R_4$$  represents a hydrogen or halogen atom; or a radical  $NR_SR_6,~S\left(O\right)_mR_7,~C\left(O\right)R_7,~alkyl,~haloalkyl or <math display="inline">OR_8$  or a radical  $-N=C\left(R_0\right)\left(R_{10}\right);$ 

 $R_5$  and  $R_6$  independently represent a hydrogen atom or an alkyl, haloalkyl, C(0) alkyl or S(0)  $_{\rm r}CF_3$  radical; or  $R_5$  and  $R_6$  may together; form a divalent alkylene radical which may be interrupted by one or two divalent heteroratoms, such as exygeneor sulphur:

 $$R_7$$  represents an alkyl or haloalkyl radical;  $$R_8$$  represents an alkyl or haloalkyl radical or a hydrogen atom;

 $$\rm R_{9}$$  represents an alkyl radical or a hydrogen atom;

 $R_{10}$  represents a phenyl or heteroaryl group optionally substituted with one or more halogen atoms or groups such as OH, -O-alkyl, -S-alkyl, cyano or alkyl;

 $$R_{11}$$  and  $$R_{12}$$  represent, independently of each other, a hydrogen or halogen atom;

 $$R_{13}$$  represents a halogen atom or a haloalkyl, haloalkoxy, S(O)  $_0CF_3$  or SF, group;

m, n, q and r represent, independently of each other, an integer equal to 0, 1 or 2;

 $\,$  X represents a trivalent nitrogen atom or a radical C-R<sub>12</sub>, the other three valency positions of the carbon atom forming part of the aromatic ring,

with the proviso that when  $R_1$  is methyl, then  $R_3$  is haloalkyl,  $R_4$  is  $NH_2,\ R_{11}$  is Cl,  $R_{13}$  is CF $_3$  and X is N .

Compounds of formula (I) in which  $R_1$  is CN will be selected most particularly. The compounds in which  $R_2$  is  $S(O)_n R_3$ , preferably with n=1,  $R_3$  preferably being CF<sub>3</sub> or alkyl, for example methyl or ethyl, or alternatively n=0,  $R_3$  preferably being CF<sub>3</sub>, as well as

10

15

20

25

30

those in which  $X = C-R_{12}$ ,  $R_{12}$  being a halogen atom, will also be selected. Compounds in which  $R_{11}$  is a halogen atom and those in which  $R_{13}$  is haloalkyl, preferably  $CF_{3}$ , are also preferred. In the context of the present invention, compounds combining two or more of these characteristics will advantageously be selected.

A preferred class of compounds of formula (I) consists of compounds such that  $-R_1$  is CN,  $-R_3$  is halo—alkyl, preferably CF<sub>3</sub>, or ethyl,  $R_4$  is NH<sub>2</sub>,  $R_{11}$  and  $R_{12}$  are, independently of each other, a halogen atom, and/or  $R_{13}$  is haloalkyl.

In the present invention, the alkyl radicals may contain generally from 1 to 6 carbon atoms. The cycle formed by the divalent alkylene radical representing  $R_{\text{S}}$  and  $R_{\text{G}}$ , as well as with the nitrogen atom to which  $R_{\text{S}}$  and  $R_{\text{G}}$  are linked may be generally a cycle of 5, 6 or 7 links.

A compound of formula (I) which is most particularly preferred in the invention is

 $1-[2,6-Cl_24-CF_3pheny1]\,3-CN\ 4-[SO-CF_3]\,5-NH_2py-razole,$ 

which is referred to hereinbelow as compound A.

Mention may also be made of the two compounds which differ from the above A by the following characteristics:

1- n = 0, 
$$R_3 = CF_3$$
  
2- n = 1,  $R_3 = ethvl.$ 

Compounds of formula (I) may be prepared according to one or other of the processes described in patent applications WO-A-87/3781, 93/6089, 94/21606 or European patent application EP-A-0,295,117, or any other process falling within the competence of a specialist skilled in the art of chemical synthesis. For the

10

15

25

30

chemical preparation of the products of the invention, a person skilled in the art is considered as having at his disposal, inter alia, all of the contents of "Chemical Abstracts" and the documents which are cited therein.

However, low concentrations, of from 1 to 15% by weight and more particularly, especially for compound A, of from 1.25 to 10%; are preferred.

Under optimum conditions, compound (I) and especially compound A is present in the collar in a proportion of from 2 to 6% by weight, more particularly from 2.5 to 5% by weight.

It would obviously be possible to add to compound A any other insecticide which might be considered useful.

These insecticides may be present in the same matrix as the compound according to the invention. A composite collar made of at least two parts, each including a different active substance, may also be used.

Within the scope of the invention, matrices usually used to make collars may be used. Preferred examples which may be mentioned are matrices based on PVC (polyvinyl chloride), as described in US-A-3,318,769, 3,852,416 and 4,150,109 and 5,437,869, and other vinyl polymers.

The plasticizers may be chosen in particular from adipates, phthalates, phosphates and citrates.

One or more plasticizers will preferably be added to the PVC, these plasticizers being chosen in particular from the following compounds:

- diethyl phthalate
- dioctyl sebacate

10

15

25

30

- dioctyl adipate
- -diisodecyl phthalate
- acetvl tributvl citrate
- diethyl hexyl phthalate
- di-n-butyl phthalate
  - benzyl butyl phthalate
  - acetyl tributyl citrate . ..
  - tricresyl phosphate
  - 2-ethylhexyl diphenyl phosphate.

Even more preferably, a PVC matrix will be used in the presence of a primary remanent plasticizer and a secondary plasticizer, in particular according to EP-A-0.539.295 and EP-A-0.537.998.

Among the secondary plasticizers, mention may be made of the following products:

- acetyl triethyl citrate
- triethyl citrate
- triacetin
- diethylene glycol monoethyl ether
- 20 triphenyl phosphate.

 $\label{eq:Acommon stabilizer may} \mbox{ also be added} \\ \mbox{thereto.}$ 

For the purposes of the present invention, the term external device should be understood to refer to any device which can be attached externally to the animal in order to provide the same function as a collar.

By varying the concentration and/or composition of the matrix, collars or other external devices according to the invention, which ensure effective and long-lasting protection against fleas, may be made. Collars or other external devices may be made with an efficacy of greater than 6 months, in particular

15

20

25

30

of greater than or equal to 12 or 18 months, even when the collar or external device is taken off for a relatively prolonged period. When the collar or device is taken off, the duration of effective protection may range from 2 to 3 months.

By varying the concentration and/or composition of the matrix, it is possible to make collars or other external devices according to the invention which ensure effective and long-lasting protection against ticks. Collars or external devices may be made with an efficacy of greater than 3 months, in particular of greater than or equal to 12 or 15 months, even when the collar or external device is taken off for a relatively prolonged period. When the collar or external device is taken off, the duration of effective protection may range from 1 to 2 months.

It is noteworthy that this very long-lasting and total efficacy is obtained by the compound according to the invention alone, without addition of another insecticide.

The subject of the present invention is also a method for eliminating ectoparasites, in particular fleas and ticks, from pets such as cats and dogs, in which method at least one collar or other external device in accordance with the invention is attached to the animal and the animal is afforded long-lasting, effective protection against these parasites, even when the collar or other external device is taken off. The method is capable of ensuring prevention and treating fleas and ticks to a high degree of efficacy and over a period exceeding 6 months against fleas and 3 months against ticks, the efficacy preferably being maintained over several weeks even if the collar or external device

1.0

15

20

25

30

is taken off or if there is a variation in the release of the compound (I) by the collar or external device. The indications of duration have been given above. Preferably, it is recommended that, in accordance with the method according to the invention, after it has been put on for the first time, the collar or external device should remain on the animal for at-least 24 hours in order for sufficient active substance to pass to the animal and for the sebaceous glands to have been able to store this active substance.

The object of this method is non-therapeutic and in particular relates to the cleaning of animal hairs and skin by elimination of the parasites which are present, as well as their residues and dejections. The treated animals thus have hair which is more pleasant to look at and to feel.

The invention also relates to such a method for therapeutic purposes, intended to treat and prevent parasitoses having pathogenic consequences.

The subject of the present invention is also the use of a compound corresponding to formula I for the production of a collar or other external device intended to be attached to a pet, in particular cats and dogs, this compound being capable of ensuring prevention and treating fleas and ticks to a high degree of efficacy and over a period exceeding 6 months against fleas and 3 months against ticks, the efficacy preferably being maintained for several weeks if the collar or the external device is taken off or lost or if there is a variation in the release of the compound (I) by the collar or the external device. The devices as above are concerned.

In particular, the method and device accor-

15

20

3

ding to the invention are designed to provide for an efficacy of greater than 95%, or even greater than 98 or 99% against fleas.

For ticks, the desired efficacy exceeds 80% or 90%.

Similarly, the invention, against fleas, is directed towards the production of a long-lasting efficacy, longer than or equal to 12 months and even to 18 months.

For ticks, this duration is longer than 12 months, or even longer than 15 months.

Also preferably, the invention is directed towards the production of collars or other external devices which make it possible to obtain an efficacy maintained in the absence of the collar or external device over a period ranging from 2 to 3 months, or more, against fleas and from 1 to 2 months, or more, against ticks.

The present invention will now be described in greater detail with the aid of non-limiting examples from which other particular features and advantages of the invention will emerge.

#### EXAMPLE 1:

#### Collar containing 10% compound A

25 The following two types of collar were prepared (mixing and then extrusion):

	(mixing and	ther	ext	rus	ior	1)	:						
	Formulation	1:	PVC										50.0%
			Stab	ili	zeı	c							0.5%
			Epox	idi	zec	i i	so	ybe	ean	0	il		5.0%
30			Diis	000	ty.	1 .	ad	ipa	ate				34.5%
			Comp	oun	d 1	Ą							10.0%
	Formulation	2:	PVC										50.0%
			Stab	ili	ze:	r							0.5%

20

5 (% by weight).

For the tests, 9 adult dogs which had not received any insecticide or acaricide for at least 40 days were chosen. The dogs were washed with an insecticide-free shampoo and combed in order to remove any existing parasites.

The dogs were divided into groups of three.

#### Group A:

Untreated controls

#### Group B:

15 Collar containing 10% compound A - formulation 1.
Group C:

Collar containing 10% compound A - formulation 2.

The dogs are infested with about 100  $\pm$  10 Ctenocephalides felis cat fleas (Unfed cat fleas) and 50  $\pm$  2 Rhipicephalus sanguineus ticks (Brown dog tick).

The treatment follows the following general schema:

- -2 Infestation with the fleas and ticks.
- O Collars are put on.
- 25 2 Fleas and ticks are counted with a comb.
  - 7 Infestation with ticks.
  - 8 Infestation with fleas.
  - 9 Counting with a comb
  - 35 Infestation with ticks.
- 30 36 Infestation with fleas.
  - 37 Counting with a comb.
  - 63 Infestation with ticks.
  - 64 Infestation with fleas.

65 Counting with a comb.

The same process is continued with infestation with fleas and ticks every month for as long as a satisfactory efficacy is observed.

Hair samples were taken on:

D 205 and D 261 for all the dogs;

The collars were taken off on D.149 on dogs 363 and 289, on D 177 for dogs 87 and 300 and on D 205 for dogs  $^{256}$  and  $^{335}$ .

The concentrations of active substance according to the invention on the hairs which were obtained at D 205 and D 261 are presented in Tables 1 and 2 respectively, in  $\mu g/g$  of hair.

<u>Table 1:</u> Concentration of active substance according to the invention on D 205.

DOG		CONCENT	RATION (µ	g.g <sup>-1</sup> )	
	UNDER THE	RIGHT	LEFT	MIDDLE	LUMBAR
No.	COLLAR	FLANK	FLANK	OF BACK	REGION
GROUP B					
363	18.63	3.11	2.33	1.73	6.99
87	19.03	2.28	3.08	5.81	4.54
256	14.66	3.04	3.01	2.48	10.99
GROUP C	·				
289	27.47	11.75	15.19	8.99	30.9
300	482.4	34.6	15.69	17.97	36.7
335	257.4	23.86	28.86	22.8	41.2

<u>Table 2:</u> Concentration of active substance according to the invention, on hairs on D 261.

1.5

20

25

30

DOG		CONCENT	RATION (μ	g.g <sup>-1</sup> )	
	UNDER THE	RIGHT	LEFT	MIDDLE	LUMBAR
No.	COLLAR	FLANK	FLANK	OF BACK	REGION
GROUP B					
363	0.94	0.65	>0.29	>0.39	1.47
87	0.64	0.94	>0.32	>0.40	ND
256	>0.97	ND	0.81	." 0.78	0.69
GROUP C	ľ				
289	3.16	>0.75	>0.59	>0.60	2.03
300	3.79	>0.70	2.04	1.35	5.52
335	1.84	1.67	2.99	4.61	4.19

#### ND = Not determined.

A distribution of active substance over the hair is observed: the concentrations on the different areas are higher with the formulation of Group C and are relatively homogeneous from one area to another.

The concentrations of active substance are still detectable 16 weeks after the collar has been taken off and are still effective at that stage.

Tables 3, 4 and 5 make it possible to carry out a concentration/activity correlation.

20

5

10

Table 3: Concentration/activity relationship on D 149.

	DOG	CONCENTRAT:	ION (μg		ACTIVIT	Y (1)	
		UNDER THE	BACK	BACK AND FLANKS			TICKS
5	No.	COLLAR	AV.	MIN	MAX.		
	GROUP B	559.5	15: 6	10:7	24 8	0	- eq
10	GROUP C	339.3	13.0	10.7	24.0	0	ī
10	289	871.3	48.9	30.8	94.4	0	00

(1) Number of fleas and ticks present on the dog

Table 4: Concentration/activity relationship on D 205.

DOG	CONCENTRAT	ION (μg	ON (μg.g <sup>-1</sup> )			ACTIVITY(1)		
	UNDER THE	BACK	AND FL	FLEAS	TICKS			
No.	COLLAR	AV.	MIN	MAX				
GROUP								
В								
363	18.6	3.54	1.73	6.99	0	12		
87	19.0	3.93	2.28	5.81	0	22		
256	15.0	4.88	2.48	11.0	0	15		
GROUP								
C								
289	27.5	16.7	8.99	30.9	0	12		
300	482.4	26.3	15.7	36.7	0	0		
335	257.4	29.2	22.8	41.2	0	0		

(1): Number of fleas and ticks present on the dog

15

20

25

Table 5: Concentration/activity relationship on D 261.

	DOG	CONCENTRAT:	ION (µg	.g <sup>-1</sup> )		ACTIVITY(1)		
		UNDER THE	BACK	BACK AND FLANKS			TICKS	
5	No.	COLLAR	AV.	MIN	MAX.			
LO	GROUP B 363 87 256	0.94 0.64 >0.97		>0.29 <loq <loq< td=""><td>1.47 0.94 0.81</td><td>0 1</td><td>21 12 24</td></loq<></loq 	1.47 0.94 0.81	0 1	21 12 24	
15	GROUP C 289 300 335	3.16 3.79 1.84		>0.59 >0.70 1.67	2.03 5.52 4.61	0 0 0	28 19 3	

(1): Number of fleas and ticks present on the dog

AV. = average

MIN = minimum

20 MAX = maximum

LOQ = limit of quantification = 0.25  $\mu$ g/g

On D 261, the activity towards fleas is 100% on all the dogs, except for one dog (presence of a flea).

The minimum concentrations (on the hair) which are 25 effective towards fleas and ticks are determined to be about:

- 20 micrograms per gram of hair for ticks
- 1 microgram per gram of hair for fleas.

An activity of at least 5 months for ticks and of at least 9 months for fleas is observed with these formulations.

#### EXAMPLE 2:

Dogs were divided up into 6 groups of 8 dogs:

35 - A: Control group: collar containing no active

substance.

- B: Reference product: commercial collar against fleas and ticks, containing 8% Chlorpyrifos (0,0-diethyl-0-(3,5,6-trichloro-2-pyridyl)phosphorothioate), referred
- 5 to as ref. in the tables.
  - C: Collar containing 2.5% compound A.
  - D: Collar containing 5% compound A. . .
  - E: Collar containing 10% compound A
- The collars were made with the same ingredients as in formulation 2 of Example 1, with, in addition, a pigment (titanium dioxide).

The tests follow the following schedule:

- 2 Infestation with fleas and ticks
- 0 Collars are put on
- 15 2 Counting with a comb
  - 7 Reinfestation with ticks
  - 8 Reinfestation with fleas
  - 9 Counting with a comb
  - 35 Reinfestation with ticks
- 20 36 Reinfestation with fleas
  - 37 Counting with a comb

The process is continued along these lines from  $\ensuremath{\mathsf{month}}$  to  $\ensuremath{\mathsf{month}}$  .

Infestations and reinfestations are carried out at a rate of:

100 ± 10 fleas

50 ± 3 ticks

Tables 6 and 7 give the average effectiveness results.

30 <u>Table 6:</u> Percentage effectiveness on fleas, calculated as a geometric mean

Ç.
1
F
J
O
36
100
(5
100
ų,
40
O

			GROUPS	TREATED	
	DAY	Ref.	2.5%	5%	10%
		8%	<u> </u>		
	-5	0.0%	9.3%	0.0%	0.0%
	2	63.9%	98.7%	100.0%	100.0%
5	9	96.3%	100.0%	100.0%	100.0%
	37	96.1%	99.9%	100.0%	100.0%
	65	99.1%	100.0%	100.0%	100.0%
	93	99.6%	100.0%	100.0%	100.0%
	121	99.4%	100.0%	100.0%	100.0%
10	149	98.3%	100.0%	100.0%	100.0%
	177	97.9%	100.0%	100.0%	100.0%
	205	98.5%	100.0%	100.0%	100.0%
	233	96.6%	100.0%	100.0%	100.0%
	268	89.6%	100.0%	100.0%	100.0%
15	289	68.4%	100.0%	100.0%	100.0%
	317	76.0%	100.0%	100.0%	100.0%
	345	79.5%	100.0%	100.0%	100.0%
	373	-	100.0%	100.0%	100.0%
	401	-	99.4%	100.0%	99.8%
20	429	-	99.2%	99.8%	99.3%
	457	-	96.9%	98.8%	99.2%
	485	-	96.7%	98.2%	98.7%
	513 ·	-	90.3%	99.0%	99.3%
25	541 .	l		98.8%	100.0%

<u>Table 7:</u> Percentage effectiveness on ticks, calculated as a geometric mean

		r			
			GROUPS	TREATED	
5	DAY	Ref.	2.5%	5%	10%
		8%			
	-5	4.7%	19.1%	19.1%	32.0%
	2	34.5%	67.0%	100.0%	98.5%
	9	86.9%	96.2%	100.0%	100.0%
	37	0.0%	27.2%	95.4%	95.4%
10	65	31.7%	85.0%	100.0%	95.8%
	93	31.8%	86.7%	100.0%	95.7%
	121	65.2%	91.3%	88.8%	92.1%
	149	59.0%	93.4%	92.0%	94.5%
	177	67.2%	84.9%	95.3%	95.2%
15	205	62.0%	86.2%	100.0%	98.4%
	233	62.6%	89.8%	98.3%	97.9%
	268	57.5%	77.4%	98.2%	99.5%
	289	44.6%	92.4%	97.2%	95:1%
	317	-	74.5%	85.6%	92.4%
20	345	-	88.1%	98.8%	98.0%
	373	-	72.4%	93.9%	95.7%
	401	-	63.4%	98.5%	93.6%
	429	-	75.3%	83.2%	93.3%
	457	-	62.4%	84.1%	86.7%
25	485	-	-	-	-
	513	-	-	-	-
	541	-		_	-

#### CLAIMS

1. Anti-flea and anti-tick collar or other external device for a pet, in particular a cat or dog, made of a matrix in which is incorporated from 0.1 to 40% by weight, relative to the collar, of a substance which is active against fleas and ticks, this active substance being formed of at least one compound corresponding to formula (I) below:

10

15

30

20 in which:

R<sub>1</sub> is CN or methyl or a halogen atom;

 $\rm R_2$  is  $\rm S\left(0\right)_n R_3$  or 4,5-dicyanoimidazol-2-yl or haloal-kyl;

R<sub>3</sub> is alkyl or haloalkyl;

or  $OR_8$  or a radical  $-N=C(R_9)(R_{10})$ ;

 $R_{s}$  and  $R_{6}$  independently represent a hydrogen atom or an alkyl, haloalkyl, C(O)alkyl, alkoxycarbonyl or S(O) $_{t}$ CF $_{3}$  radical; or  $R_{5}$  and  $R_{6}$  may together form a divalent alkylene radical which may be interrupted by

one or two divalent hetero atoms, such as oxygen or sulphur;

10

15

20

25

30

R, represents an alkyl or haloalkyl radical;

 $\ensuremath{R_{8}}$  represents an alkyl or haloalkyl radical or a hydrogen atom;

 $R_9$  represents an alkyl radical or a hydrogen atom;  $R_{10} \mbox{ represents a phenyl or heteroaryl group optionally substituted with one or more halogen atoms or groups such as OH, -O-alkyl, -S-alkyl, cyano or alkyl;$ 

 $R_{11}$  and  $R_{12}$  represent, independently of each other, a hydrogen or halogen atom, or CN or  $NO_2; \\$ 

 $R_{13}$  represents a halogen atom or a haloalkyl, haloalkoxy,  $S(0)_q C F_3$  or  $S F_5$  group;

m, n, q and r represent, independently of each other, an integer equal to 0, 1 or 2;

X represents a trivalent nitrogen atom or a radical  $C-R_{12}$ , the other three valency positions of the carbon atom forming part of the aromatic ring;

with the proviso that when  $R_1$  is methyl, either  $R_3$  is haloalkyl,  $R_4$  is NH<sub>2</sub>,  $R_{11}$  is Cl,  $R_{13}$  is CF<sub>3</sub> and X is N; or  $R_2$  is 4,5-dicyanoimidazol-2-yl,  $R_4$  is Cl,  $R_{11}$  is Cl,  $R_{13}$  is CF<sub>3</sub> and X is =C-Cl;

this collar or other external device being designed to ensure more than 6 months of efficacy against fleas and more than 3 months of efficacy against ticks, the efficacy preferably being maintained for several weeks even if the collar or other external device is taken off or lost or if there is a variation in the release of the compound (I) by the matrix.

2. Collar according to claim 1, characterized in that the compound of formula (I) is such that:

R, is CN or methyl;

R<sub>2</sub> is S(O)<sub>2</sub>R<sub>3</sub>;

R3 is alkyl or haloalkyl;

R, represents a hydrogen or halogen atom; or a

10

15

20

25

radical  $NR_5R_6$ ,  $S(O)_mR_7$ ,  $C(O)R_7$ , alkyl, haloalkyl or  $OR_8$  or a radical  $-N=C(R_9)$   $(R_{10})$ ;

 $R_{\rm S}$  and  $R_{\rm G}$  independently represent a hydrogen atom or an alkyl, haloalkyl, C(O)alkyl or S(O) $_{\rm r}CF_{\rm 3}$  radical; or  $R_{\rm S}$  and  $R_{\rm G}$  may together form a divalent alkylene radical which may be interrupted by one or two divalent hetero atoms, such as oxygen or sulphur;

 $R_7$  represents an alkyl or haloalkyl radical;

 $R_{8}$  represents an alkyl or haloalkyl radical or a hydrogen atom;

 $R_{9}$  represents an alkyl radical or a hydrogen atom;  $R_{10} \text{ represents a phenyl or heteroaryl group optionally substituted with one or more halogen atoms or groups such as OH, -O-alkyl, -S-alkyl, cyano or alkyl; } \\$ 

 $R_{11}$  and  $R_{12}$  represent, independently of each other, a hydrogen or halogen atom;

 $R_{13}$  represents a halogen atom or a haloalkyl, haloalkoxy,  $S(0)_{\sigma}CF_3$  or  $SF_9$  group;

m, n, q and r represent, independently of each other, an integer equal to 0, 1 or 2;

X represents a trivalent nitrogen atom or a radical C-R $_{12}$ , the other three valency positions of the carbon atom forming part of the aromatic ring;

with the proviso that when  $R_1$  is methyl, then  $R_3$  is haloalkyl,  $R_4$  is  $NH_2,\ R_{11}$  is Cl,  $R_{13}$  is CF, and X is N .

- 3. Coilar according to claim 2, wherein the compound of formula (I) is such that  $R_1$  is CN.
- 4. Collar according to claim 2, wherein the compound of formula (I) is such that  $R_{13}$  is haloalkyl.
- 30 5. Collar according to claim 4, wherein the compound of formula (I) is such that  $R_{13}$  is  $CF_3$ .
  - 6. Collar according to claim 2, wherein the compound of formula (I) is such that  $R_2$  is  $S(0)_nR_3$ .

- 7. Collar according to claim 6, wherein n=1 and  $R_3$  is chosen among the group consisting of  $CF_{3}$ , methyl, ethyl.
- 8. Collar according to claim 6, wherein n=0 and  $R_3$  is CF,
- 9. Collar according to claim 2,wherein the compound of formula (I) is such that X is  $C-R_{12},\ R_{12}$  being a halogen atom.
- 10. Collar according to claim 2, wherein the compound of formula (I) is chosen from those in which  $R_1$  is
- CN,  $R_3$  is haloalkyl,  $R_4$  is  $NH_2$ ,  $R_{11}$  and  $R_{12}$  are, independently of each other, a halogen atom, and/or  $R_{13}$  is haloalkyl.
- 11. Collar according to claim 2, wherein the compound of formula (I) is chosen among the group consisting of compound A:
  - l-[  $^{\prime}2,6\text{-Cl}_24\text{-CF}_3\text{phenyl}]3\text{-CN4-[SO-CF}_3]5\text{-NH}_2\text{pyrazole}$  and its derivatives with n=0 and  $R_3$  is CF3, and n=1 and  $R_3$  is ethyl.
- 20 12. Collar according to claim 2, wherein the collar comprises from 1 to 15% active substance.
  - 12. Collar according to claim 2, wherein the collar comprises from 1.25 to 10% active substance.
  - 13. Collar according to claim 2, wherein the collar comprises from 2 to 6% active substance.
- 25 comprises from 2 to 6% active substance.

  14. Collar according to claim 2 wherein
  - 14. Collar according to claim 2, wherein the collar comprises from 2.5 to 5% active substance.
  - 15. Collar according to claim 11, wherein the collar comprises from 1.25 to 10% active substance.
- 30 16. Collar according to claim 11, wherein the collar comprises from 2 to 6% active substance.
  - 17. Collar according to claim 11, wherein the collar comprises from 2.5 to 5% active substance.

18. Collar according to claim 11, wherein the efficacy is maintained when the collar or external device is taken off or lost, over a period ranging from 2 to 3 months against fleas and from 1 to 2 months against ticks.

19. Collar according to claim 11, wherein it comprises a concentration of active substance which ensures effective protection against fleas for a period longer than or equal to 12 or 18 months.

20. Collar according to claim 11, wherein it comprises a concentration of active substance which ensures effective protection against ticks for a period longer than or equal to 12 or 15 months.

21. Method for eliminating fleas and ticks from pets, in particular cats and dogs, in which one attaches to the pets at least one collar or other external device which comprises a compound corresponding to formula (I) below:

20

15

5

25

30 in which:

 $\rm R_1$  is CN or methyl or a halogen atom;  $\rm R_2$  is S(O)  $_n R_3$  or 4,5-dicyanoimidazol-2-yl or haloal-kyl;

10

15

20

25

30

R, is alkyl or haloalkyl;

 $R_4$  represents a hydrogen or halogen atom; or a radical  $NR_5R_6$ ,  $S(0)_mR_7$ ,  $C(0)R_7$ ,  $C(0)O-R_7$ , alkyl, haloalkyl or  $OR_0$  or a radical  $-N=C(R_9)(R_{10})$ ;

 $R_{\delta}$  and  $R_{\delta}$  independently represent a hydrogen atom or an alkyl, haloalkyl,  $C\left(0\right)$  alkyl, alkoxycarbonyl or  $S\left(0\right)_{\tau}CF_{3}$  radical; or  $R_{\delta}$  and  $R_{\delta}$  may together form a divalent alkylene radical which may be interrupted by one or two divalent hetero atoms, such as oxygen or sulphur;

R, represents an alkyl or haloalkyl radical;

 $\ensuremath{R_{\text{R}}}$  represents an alkyl or haloalkyl radical or a hydrogen atom;

 $\rm R_9$  represents an alkyl radical or a hydrogen atom;  $\rm R_{10}$  represents a phenyl or heteroaryl group optionally substituted with one or more halogen atoms or groups such as OH, -O-alkyl, -S-alkyl, cyano or alkyl;

 $R_{11}$  and  $R_{12}$  represent, independently of each other, a hydrogen or halogen atom, or optionally CN or  $NO_2\,;$ 

 $R_{13}$  represents a halogen atom or a haloalkyl, haloalkoxy,  $S(0)\,_qCF_3$  or  $SF_5$  group;

m, n, q and r represent, independently of each other, an integer equal to 0, 1 or 2;

X represents a trivalent nitrogen atom or a radical C-R $_{12}$ , the other three valency positions of the carbonatom forming part of the aromatic ring;

with the proviso that when  $R_1$  is methyl, either  $R_3$  is haloalkyl,  $R_4$  is NH<sub>2</sub>,  $R_{11}$  is Cl,  $R_{13}$  is CF<sub>3</sub> and X is N; or  $R_2$  is 4,5-dicyanoimidazol-2-yl,  $R_4$  is Cl,  $R_{11}$  is Cl,  $R_{13}$  is CF<sub>3</sub> and X is =C-Cl;

which method ensuring prevention and treating fleas and ticks to a high degree of efficacy and over a period exceeding 6 months against fleas and 3 months against ticks, the efficacy preferably being maintained over

several weeks even if the collar or external device is taken off or if there is a variation in the release of the compound (I) by the collar or external device.

22. Method according to claim 21, wherein the compound of formula (I) is such that:

R, is CN or methyl;

R, is S(O),R3;

R, is alkyl or haloalkyl;

 $R_4$  represents a hydrogen or halogen atom; or a radical NRsR6, S(O)\_mR7, C(O)R7, alkyl, haloalkyl or OR8 or a radical -N=C(R9) (R10);

 $R_{\text{S}}$  and  $R_{\text{G}}$  independently represent a hydrogen atom or an alkyl, haloalkyl, C(0)alkyl or S(0) $_{\text{r}}\text{CF}_{3}$  radical; or  $R_{\text{S}}$  and  $R_{\text{G}}$  may together form a divalent alkylene radical which may be interrupted by one or two divalent hetero atoms, such as oxygen or sulphur;

R, represents an alkyl or haloalkyl radical;

 $R_{\theta}$  represents an alkyl or haloalkyl radical or a hydrogen atom;

 $$\rm R_9$$  represents an alkyl radical or a hydrogen atom;  $$\rm R_{10}$$  represents a phenyl or heteroaryl group optionally substituted with one or more halogen atoms or groups such as OH, -O-alkyl, -S-alkyl, cyano or alkyl;

 $\underset{i=1}{R}_{11}$  and  $R_{12}$  represent, independently of each other, a hydrogen or halogen atom;

 $\rm R_{13}$  represents a halogen atom or a haloalkyl, haloalkoxy, S(O)  $_{\rm q}{\rm CF_3}$  or SF, group;

m, n, q and r represent, independently of each other, an integer equal to 0, 1 or 2;

X represents a trivalent nitrogen atom or a radical C-R $_{12}$ , the other three valency positions of the carbon atom forming part of the aromatic ring;

with the proviso that when  $R_{\text{1}}$  is methyl, then  $R_{\text{3}}$  is

20

2.5

30

5

10

20

30

haloalkyl,  $R_4$  is  $NH_2$ ,  $R_{11}$  is Cl,  $R_{13}$  is  $CF_3$  and X is N.

- 23. Method according to claim 22, wherein the compound of formula (I) is such that  $R_1$  is CN.
- 24. Method according to claim 22, wherein the compound of formula (I) is such that  $R_{13}$  is haloalkyl.
- 25. Method according to claim 22, wherein the compound of formula (I) is such that  $R_{13}$ , is  $CF_3$ .
- 26. Method according to claims 22, wherein the compound of formula (I) is such that  $R_2$  is  $S(0)_n R_3$ .
- 10 27. Method according to claim 26, wherein n=1 and  $R_3$  is chosen among the group consisting of  $CF_3$ , methyl, ethyl.
  - 28. Method according to claim 26, wherein n = 0 and  $R_{\rm 3}$  is CF<sub>3</sub>.
- 15 29. Method according to claim 22, wherein the compound of formula (I) is such that X is  $C-R_{12}$ ,  $R_{12}$  being a halogen atom.
  - 30. Method according to claim 22, wherein the compound of formula (I) is such that  $R_1$  is CN,  $R_3$  is haloal-kyl,  $R_4$  is NH<sub>2</sub>,  $R_{11}$  and  $R_{12}$  are, independently of each other, a halogen atom, and/or  $R_{13}$  is haloalkyl.
  - 31. Method according to claim 22, wherein the compound of formula (I) is chosen among the group consisting of compound A:
- - 32. Method according to claim 22, wherein the compound of formula (I) is present in a proportion of from 1 to 15% by weight.
    - 33. Method according to claim 22, wherein the compound of formula (I) is present in a proportion of from 1.25 to 10%.

15

- 34. Method according to claim 22, wherein the compound of formula (I) is present in a proportion of from 2 to 6%.
- 35. Method according to claim 22, wherein the compound of formula (I) is present in a proportion of from 2.5 to 5% by weight.
  - 36. Method according to claim 31, wherein the compound of formula (I) is present in a proportion of from 1.25 to 10%.
- 10 37. Method according to claim 31, wherein the compound of formula (I) is present in a proportion of from 2 to 6% by weight.
  - 38. Method according to claim 31, wherein the compound of formula (I) is present in a proportion of from 2.5 to 5% by weight.
  - 39. Method according to claim 31, wherein the efficacy is greater than 95% against fleas.
  - 40. Method according to claim 31, wherein the efficacy is greater than 98% or 99% against fleas.
  - 41. Method according to claim 31, wherein the efficacy is greater than 80% against ticks.
    - 42. Method according to claim 31, wherein the efficacy is greater than 90% against ticks.
- 43. Method according to claim 31, wherein the long-25 lasting efficacy is longer than or equal to 12 months against fleas.
  - 44. Method according to claim 31, wherein the longlasting efficacy is longer than or equal to 18 months against fleas.
- 30 45. Method according to claim 31, wherein the longlasting efficacy is longer than or equal to 12 months against ticks.
  - 46. Method according to claim 31, wherein the long-

lasting efficacy is longer than or equal to  $15\ \mathrm{months}$  against ticks.

47. Method according to claim 31, wherein the efficacy is maintained when the collar or external device is taken off or lost, over a period ranging from 2 to 3 months against fleas and from 1 to  $^{\circ}2$  months against ticks.

#### COMPANY KNOWN AS: RHONE MERIEUX

N-Phenylpyrazole-based anti-flea and anti-tick collar for cats and dogs

#### ABSTRACT

Collar or other external device for a pet, in particular a cat or dog, made of a matrix in which is incorporated from 0.1 to 40% by weight, preferably from 1 to 15 % by weight, relative to the collar, of a substance which is active against ectoparasites such as fleas and ticks, this active substance being formed of at least one compound corresponding to formula (I) below:

this collar or other external device being designed to ensure more than 6 months of efficacy against fleas and more than 3 months of efficacy against ticks, the efficacy being maintained for several weeks even if the collar or other external device is taken off or lost or if there is a variation in the release of the compound.

# DECLAR TON FOR USA PATENT APPLIC TION (including Design and National Stage PCT) Attorney's Docket ID: XI/P03139US1

(including Design and National Stage PCT)

Attorney's Docket ID: XI/P03139US

As a below named inventor, I hereby deel My residence, post office address and citizens s listed below) or an original, first and joint is on the invention entitled	inventor (if plural names are listed be	y name. I believe I am the original, first and so low) of the subject matter which is claimed and LEA AND ANTI-TICK COLLAR FO	le inventor (if only one name for which a patent is sought DR CATS AND DOGS
		, the s	pecification of which
is attached hereto. (or)			
x was filed on May 27, 1997	, [ ] and was amended on		
[ X] as U.S. Application No(	08/863,182	or	
[ ] as International PCT Applica	ation No.		
hereby state that I have reviewed and underso above. I acknowledge the duty to disclost hereby claim foreign priority benefits under settificate or \$365 (a) of any PCT Internation	tand the contents of the above-identifies information which is material to pair.  Title 35, United States Code, § II: all application which designated at lea	ed specification, including the claims, as amend- tentability as defined in Title 37, Code of Fede 9 (a) - (d) or §365 (b) of any foreign applicati st one country other than the United States of A t or inventor's certificate, or any PCT Internation	ed by any amendment referred ral Regulations, § 1.56. on(s) for patent or inventor': merica, listed below and have
also identified below, where priority is not clar late before that of the application on which	imed, any foreign application for pater priority is claimed:	at or inventor's certificate, or any PCT Internation	nal application, having a filing
rior Foreign Application(s) (ADDI	ITIONAL APPLICATIONS IDENTIF	TED ON ATTACHED SHEET):	
Number	Country	Day/Month/Year Filed	Priority Not Claimed
96 04 206	FRANCE	29 March 1996	
97 03 707	FRANCE	26 March 1997	
hereby claim the benefit under Title 35. Unithe U.S., listed below, and insofar as the subpplication in the manner provided by the first patentability as defined in Title 37, Code of PCT International filing date of this applies.	ed States Code, § 120 of any United St oject matter of each claims of this app it paragraph of Title 35, United States f Federal Regulations § 1.56 which be cation. ( ADDITIONAL APPLIC	ates application(s), or §365(c) of any PCT Intern dication is not disclosed in the prior United Sta Code, § 112, I acknowledge the duty to disclose came available between the filing date of the pri ATIONS IDENTIFIED ON ATTACHED SHE	ational application designating tes or PCT International information which is materia or application and the nationa ET.)
Application Serial No.	Day/Month/Year Filed	Status - pate	nted, pending, abandoned
08/692,430	5 August 1996	Aband	oned
*	N & TAYLOR associated with the Ci	ustomer Number provided below to prosecute t nat all correspondence be addressed to that Cus	his application and to transactomer Number.
are	CUSTOMER NUM		
Direct all telephone calls toThomas	P. Sarro	, at TEL (703) 739-4900 (Fax: 703	-739-9577)
Tiereby declare that all statements made he with further that these statements were made vander § 1001 of Title 18 of the United States	rein of my own knowledge are true as with the knowledge that willful false sta Code and that such willful false statem	nd that all statements made on information and atements and the like so made are punishable by ents may jeopardize the validity of the application	belief are believed to be tru fine or imprisonment, or both n or any patent issued thereon
III Name of Sole	PANKIN	- Citizenship	FRANCE
First Inventor Philippe J	de la Peyrette - 311		FRANCE
104 Chemin	de la Peyrette - 311	70 TOOKNEPEUTEEE, PRANCE	
Residence - City, State/Country if different from P.O. address) SIGN AND DATE HERE: Inventor's Signature:	- Kalife kamin	Date: 21	1-11-97
Full Name of Second		Citizenship	
Full Post Office Address		· · · · · · · · · · · · · · · · · · ·	
Residence - City, State/Country if different from P.O. address)			
SIGN AND DATE HERE: Inventor's Signature:		Date:	
Full Name of Third		Citizenship	
Joint Inventor, if any Full Post Office Address			
Residence - City, State/Country (if different from P.O. address)			
SIGN AND DATE HERE: Inventor's Signature:		Date:	
Full Name of Fourth		Citizenship	
Joint Inventor, if any Full Post Office Address			
Residence - City, State/Country (if different from P.O. address)			
BIGN AND		D	
DATE HERE: Inventor's Signature:		Date:	

# IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Attorney's Docket: XI/P03139US2

In re application of:

JEANNIN, Philippe

Divisional Application of

Serial No. 08/863,182 : Art Unit: 1615

Filed: Concurrently herewith : Examiner: N. Levy

For: N-PHENYLPYRAZOLE-BASED ANTI-FLEA AND ANTI-TICK COLLAR FOR CATS

AND DOGS

# **CUSTOMER NUMBER and CHANGE OF ADDRESS**

Honorable Assistant Commissioner of Patents Washington, D. C. S.I.R.

Please change the address of LARSON & TAYLOR to:

LARSON & TAYLOR

Suite 900

1199 North Fairfax Street

Alexandria, Virginia, 22314

The address is associated with the Customer Number provided below. Direct all correspondence to the Customer Number for the above-identified patent application.

# **CUSTOMER NUMBER: 000881**

Respectfully submitted,

LARSON & TAYLOR

Date: October 19, 1998

Thomas P. Sarro

Registration No. 19,396